

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : A61K 31/42, 31/415, 31/425, 31/505, C07C 255/03, C07D 207/08, 217/18, 231/04, 263/04, 277/04, 285/12, 295/023, 413/10, 417/10</p>	A1	<p>(11) International Publication Number: <b>WO 98/53818</b></p> <p>(43) International Publication Date: 3 December 1998 (03.12.98)</p>												
<p>(21) International Application Number: PCT/US98/10952</p> <p>(22) International Filing Date: 29 May 1998 (29.05.98)</p> <p>(30) Priority Data:</p> <table border="0"> <tr> <td>60/047,954</td> <td>29 May 1997 (29.05.97)</td> <td>US</td> </tr> <tr> <td>9714335.8</td> <td>7 July 1997 (07.07.97)</td> <td>GB</td> </tr> <tr> <td>60/066,787</td> <td>25 November 1997 (25.11.97)</td> <td>US</td> </tr> <tr> <td>9800684.4</td> <td>14 January 1998 (14.01.98)</td> <td>GB</td> </tr> </table>		60/047,954	29 May 1997 (29.05.97)	US	9714335.8	7 July 1997 (07.07.97)	GB	60/066,787	25 November 1997 (25.11.97)	US	9800684.4	14 January 1998 (14.01.98)	GB	<p>(74) Common Representative: MERCK &amp; CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p> <p>(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, GW, HU, ID, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p>
60/047,954	29 May 1997 (29.05.97)	US												
9714335.8	7 July 1997 (07.07.97)	GB												
60/066,787	25 November 1997 (25.11.97)	US												
9800684.4	14 January 1998 (14.01.98)	GB												
<p>(71) Applicant (for all designated States except US): MERCK &amp; CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): DURETTE, Philippe, L. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). HAGMANN, William, K. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). MacCOSS, Malcolm [GB/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). MILLS, Sander, G. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). MUMFORD, Richard, A. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p>		<p>Published</p> <p>With international search report.</p>												
<p>(54) Title: SULFONAMIDES AS CELL ADHESION INHIBITORS</p>														
<p>(57) Abstract</p> <p>Compounds of formula (I) are antagonists of VLA-4 and/or <math>\alpha_4\beta_7</math>, and as such are useful in the inhibition or prevention of cell adhesion and cell-adhesion mediated pathologies. These compounds may be formulated into pharmaceutical compositions and are suitable for use in the treatment of asthma, allergies, inflammation, multiple sclerosis, and other inflammatory and autoimmune disorders.</p>														

TITLE OF THE INVENTION  
SULFONAMIDES AS CELL ADHESION INHIBITORS

SUMMARY OF THE INVENTION

5           The compounds of the present invention are antagonists of the VLA-4 integrin ("very late antigen-4"; CD49d/CD29; or  $\alpha 4\beta 1$ ) and/or the  $\alpha 4\beta 7$  integrin (LPAM-1 and  $\alpha 4\beta p$ ), thereby blocking the binding of VLA-4 to its various ligands, such as VCAM-1 and regions of fibronectin and/or  $\alpha 4\beta 7$  to its various ligands, such as MadCAM-1, VCAM-1 and  
10   fibronectin. Thus, these antagonists are useful in inhibiting cell adhesion processes including cell activation, migration, proliferation and differentiation. These antagonists are useful in the treatment, prevention and suppression of diseases mediated by VLA-4 and/or  $\alpha 4\beta 7$  binding and cell adhesion and activation, such as multiple sclerosis,  
15   asthma, allergic rhinitis, allergic conjunctivitis, inflammatory lung diseases, rheumatoid arthritis, septic arthritis, type I diabetes, organ transplantation, restenosis, autologous bone marrow transplantation, inflammatory sequelae of viral infections, myocarditis, inflammatory bowel disease including ulcerative colitis and Crohn's disease, certain  
20   types of toxic and immune-based nephritis, contact dermal hypersensitivity, psoriasis, tumor metastasis, and atherosclerosis.

BACKGROUND OF THE INVENTION

25           The present invention relates to sulfonamide derivatives which are useful for the inhibition and prevention of leukocyte adhesion and leukocyte adhesion-mediated pathologies. This invention also relates to compositions containing such compounds and methods of treatment using such compounds.

30           Many physiological processes require that cells come into close contact with other cells and/or extracellular matrix. Such adhesion events may be required for cell activation, migration, proliferation and differentiation. Cell-cell and cell-matrix interactions are mediated through several families of cell adhesion molecules

(CAMs) including the selectins, integrins, cadherins and immunoglobulins. CAMs play an essential role in both normal and pathophysiological processes. Therefore, the targetting of specific and relevant CAMs in certain disease conditions without interfering with normal cellular functions is essential for an effective and safe therapeutic agent that inhibits cell-cell and cell-matrix interactions.

The integrin superfamily is made up of structurally and functionally related glycoproteins consisting of  $\alpha$  and  $\beta$  heterodimeric, transmembrane receptor molecules found in various combinations on nearly every mammalian cell type. (for reviews see: E. C. Butcher, Cell, 67, 1033 (1991); T. A. Springer, Cell, 76, 301 (1994); D. Cox et al., "The Pharmacology of the Integrins." Medicinal Research Rev. 14, 195 (1994) and V. W. Engleman et al., "Cell Adhesion Integrins as Pharmaceutical Targets." in Ann. Repts. in Medicinal Chemistry, Vol. 31, J. A. Bristol, Ed.; Acad. Press, NY, 1996, p. 191).

VLA-4 ("very late antigen-4"; CD49d/CD29; or  $\alpha_4\beta_1$ ) is an integrin expressed on all leukocytes, except platelets and mature neutrophils, including dendritic cells and macrophage-like cells and is a key mediator of the cell-cell and cell-matrix interactions of these cell types (see M. E. Hemler, "VLA Proteins in the Integrin Family: Structures, Functions, and Their Role on Leukocytes." Ann. Rev. Immunol. 8, 365 (1990)). The ligands for VLA-4 include vascular cell adhesion molecule-1 (VCAM-1) and the CS-1 domain of fibronectin (FN). VCAM-1 is a member of the Ig superfamily and is expressed *in vivo* on endothelial cells at sites of inflammation. (See R. Lobb et al. "Vascular Cell Adhesion Molecule 1." in Cellular and Molecular Mechanisms of Inflammation, C. G. Cochrane and M. A. Gimbrone, Eds.; Acad. Press, San Diego, 1993, p. 151.) VCAM-1 is produced by vascular endothelial cells in response to pro-inflammatory cytokines (See A. J. H. Gearing and W. Newman, "Circulating adhesion molecules in disease." Immunol. Today, 14, 506 (1993). The CS-1 domain is a 25 amino acid sequence that arises by alternative splicing within a region of fibronectin. (For a review, see R. O. Hynes "Fibronectins.", Springer-

Velag, NY, 1990.) A role for VLA-4/CS-1 interactions in inflammatory conditions has been proposed (see M. J. Elices, "The integrin  $\alpha_4\beta_1$  (VLA-4) as a therapeutic target" in Cell Adhesion and Human Disease, Ciba Found. Symp., John Wiley & Sons, NY, 1995, p. 79).

- 5  $\alpha_4\beta_7$  (also referred to as LPAM-1 and  $\alpha_4\beta_p$ ) is an integrin expressed on leukocytes and is a key mediator of leukocyte trafficking and homing in the gastrointestinal tract (see C. M. Parker et al., Proc. Natl. Acad. Sci. USA, **89**, 1924 (1992)). The ligands for  $\alpha_4\beta_7$  include mucosal addressing cell adhesion molecule-1 (MadCAM-1) and, upon  
10 activation of  $\alpha_4\beta_7$ , VCAM-1 and fibronectin (Fn). MadCAM-1 is a member of the Ig superfamily and is expressed in vivo on endothelial cells of gut-associated mucosal tissues of the small and large intestine ("Peyer's Patches") and lactating mammary glands. (See M. J. Briskin et al., Nature, **363**, 461 (1993); A. Hamann et al., J. Immunol., **152**, 3282  
15 (1994)). MadCAM-1 can be induced in vitro by proinflammatory stimuli (See E. E. Sikorski et al. J. Immunol., **151**, 5239 (1993)). MadCAM-1 is selectively expressed at sites of lymphocyte extravasation and specifically binds to the integrin,  $\alpha_4\beta_7$ .

- Neutralizing anti- $\alpha_4$  antibodies or blocking peptides that  
20 inhibit the interaction between VLA-4 and/or  $\alpha_4\beta_7$  and their ligands have proven efficacious both prophylactically and therapeutically in several animal models of disease, including i) experimental allergic encephalomyelitis, a model of neuronal demyelination resembling multiple sclerosis (for example, see T. Yednock et al., "Prevention of  
25 experimental autoimmune encephalomyelitis by antibodies against  $\alpha_4\beta_1$  integrin." Nature, **356**, **63** (1993) and E. Keszthelyi et al., "Evidence for a prolonged role of  $\alpha_4$  integrin throughout active experimental allergic encephalomyelitis." Neurology, **47**, 1053 (1996)); ii) bronchial hyperresponsiveness in sheep and guinea pigs as models for the various  
30 phases of asthma (for example, see W. M. Abraham et al., " $\alpha_4$ -Integrins mediate antigen-induced late bronchial responses and prolonged airway hyperresponsiveness in sheep." J. Clin. Invest. **93**, 776 (1993) and A. A. Y. Milne and P. P. Piper, "Role of VLA-4 integrin in leucocyte

- recruitment and bronchial hyperresponsiveness in the guinea-pig." Eur. J. Pharmacol., **282**, 243 (1995)); iii) adjuvant-induced arthritis in rats as a model of inflammatory arthritis (see C. Barbadillo et al., "Anti-VLA-4 mAb prevents adjuvant arthritis in Lewis rats." Arthr. Rheuma. (Suppl.), **36** 95 (1993) and D. Seiffge, "Protective effects of monoclonal antibody to VLA-4 on leukocyte adhesion and course of disease in adjuvant arthritis in rats." J. Rheumatol., **23**, 12 (1996)); iv) adoptive autoimmune diabetes in the NOD mouse (see J. L. Baron et al., "The pathogenesis of adoptive murine autoimmune diabetes requires an interaction between  $\alpha_4$ -integrins and vascular cell adhesion molecule-1.", J. Clin. Invest., **93**, 1700 (1994), A. Jakubowski et al., "Vascular cell adhesion molecule-Ig fusion protein selectively targets activated  $\alpha_4$ -integrin receptors in vivo: Inhibition of autoimmune diabetes in an adoptive transfer model in nonobese diabetic mice." J. Immunol., **155**, 938 (1995), and X. D. Yang et al., "Involvement of beta 7 integrin and mucosal addressin cell adhesion molecule-1 (MadCAM-1) in the development of diabetes in nonobese diabetic mice", Diabetes, **46**, 1542 (1997)); v) cardiac allograft survival in mice as a model of organ transplantation (see M. Isobe et al., "Effect of anti-VCAM-1 and anti-VLA-4 monoclonal antibodies on cardiac allograft survival and response to soluble antigens in mice.", Transplant. Proc., **26**, 867 (1994) and S. Molossi et al., "Blockade of very late antigen-4 integrin binding to fibronectin with connecting segment-1 peptide reduces accelerated coronary arteriopathy in rabbit cardiac allografts." J. Clin. Invest., **95**, 2601 (1995)); vi) spontaneous chronic colitis in cotton-top tamarins which resembles human ulcerative colitis, a form of inflammatory bowel disease (see D. K. Podolsky et al., "Attenuation of colitis in the Cotton-top tamarin by anti- $\alpha_4$  integrin monoclonal antibody.", J. Clin. Invest., **92**, 372 (1993)); vii) contact hypersensitivity models as a model for skin allergic reactions (see T. A. Ferguson and T. S. Kupper, "Antigen-independent processes in antigen-specific immunity.", J. Immunol., **150**, 1172 (1993) and P. L. Chisholm et al., "Monoclonal antibodies to the integrin  $\alpha_4$  subunit inhibit the murine contact hypersensitivity

- response." Eur. J. Immunol., 23, 682 (1993)); viii) acute neurotoxic nephritis (see M. S. Mulligan et al., "Requirements for leukocyte adhesion molecules in nephrotoxic nephritis," J. Clin. Invest., 91, 577 (1993)); ix) tumor metastasis (for examples, see M. Edward, "Integrins and other adhesion molecules involved in melanocytic tumor progression," Curr. Opin. Oncol., 7, 185 (1995)); x) experimental autoimmune thyroiditis (see R. W. McMurray et al., "The role of  $\alpha 4$  integrin and intercellular adhesion molecule-1 (ICAM-1) in murine experimental autoimmune thyroiditis." Autoimmunity, 23, 9 (1996); and xi) ischemic tissue damage following arterial occlusion in rats (see F. Squadrito et al., "Leukocyte integrin very late antigen-4/vascular cell adhesion molecule-1 adhesion pathway in splanchnic artery occlusion shock." Eur. J. Pharmacol., 318, 153 (1996; xii) inhibition of TH2 T-cell cytokine production including IL-4 and IL-5 by VLA-4 antibodies which would attenuate allergic responses (J. Clinical Investigation 100, 3083 (1997). The primary mechanism of action of such antibodies appears to be the inhibition of lymphocyte and monocyte interactions with CAMs associated with components of the extracellular matrix, thereby limiting leukocyte migration to extravascular sites of injury or inflammation and/or limiting the priming and/or activation of leukocytes.


- There is additional evidence supporting a possible role for VLA-4 interactions in other diseases, including rheumatoid arthritis; various melanomas, carcinomas, and sarcomas; inflammatory lung disorders; acute respiratory distress syndrome (ARDS); atherosclerotic plaque formation; restenosis; uveitis and circulatory shock (for examples, see A. A. Postigo et al., "The  $\alpha 4 \beta 1$ /VCAM-1 adhesion pathway in physiology and disease," Res. Immunol., 144, 723 (1994) and J.-X. Gao and A. C. Issekutz, "Expression of VCAM-1 and VLA-4 dependent T-lymphocyte adhesion to dermal fibroblasts stimulated with proinflammatory cytokines." Immunol. 89, 375 (1996)).

At present, there is a humanized monoclonal antibody (Antegren® Athena Neurosciences/Elan ) against VLA-4 in clinical development for the treatment of "flares" associated with multiple

- sclerosis and a humanized monoclonal antibody (ACT-1®/LDP-02 LeukoSite) against  $\alpha_4\beta_7$  in clinical development for the treatment of inflammatory bowel disease. Several peptidyl antagonists of VLA-4 have been described (D. Y. Jackson et al., "Potent  $\alpha_4\beta_1$  peptide antagonists as potential anti-inflammatory agents", *J. Med. Chem.*, **40**, 3359 (1997); H. N. Shroff et al., "Small peptide inhibitors of  $\alpha_4\beta_7$  mediated MadCAM-1 adhesion to lymphocytes", *Bioorg. Med. Chem. Lett.*, **6**, 2495 (1996); US 5,510,332, WO97/03094, WO97/02289, WO96/40781, WO96/22966, WO96/20216, WO96/01644, WO96/06108, WO95/15973). There is one report of nonpeptidyl inhibitors of the ligands for  $\alpha_4$ -integrins (WO96/31206).
- There still remains a need for low molecular weight, specific inhibitors of VLA-4- and  $\alpha_4\beta_7$ -dependent cell adhesion that have improved pharmacokinetic and pharmacodynamic properties such as oral bioavailability and significant duration of action. Such compounds would prove to be useful for the treatment, prevention or suppression of various pathologies mediated by VLA-4 and  $\alpha_4\beta_7$  binding and cell adhesion and activation.

#### DETAILED DESCRIPTION OF THE INVENTION

- One aspect of the present invention provides a method for the treatment of diseases, disorders, conditions or symptoms mediated by cell adhesion in a mammal which comprises administering to said mammal an effective amount of a compound Formula I:



25

or a pharmaceutically acceptable salt thereof wherein:

- R<sup>1</sup> is
- 1) C<sub>1-10</sub>alkyl,
  - 2) C<sub>2-10</sub>alkenyl,
  - 3) C<sub>2-10</sub>alkynyl,
  - 4) Cy,

30



- 5) Cy-C<sub>1-10</sub>alkyl,
- 6) Cy-C<sub>2-10</sub>alkenyl,
- 7) Cy-C<sub>2-10</sub>alkynyl,

5 wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from R<sup>a</sup>; and Cy is optionally substituted with one to four substituents independently selected from R<sup>b</sup>;

R<sup>2</sup> and R<sup>3</sup> are independently selected from

- 1) hydrogen, or
- 10 2) a group selected from R<sup>1</sup>;

- R<sup>4</sup> is
- 1) hydrogen,
  - 2) C<sub>1-10</sub>alkyl,
  - 3) C<sub>2-10</sub>alkenyl,
  - 15 4) C<sub>2-10</sub>alkynyl,
  - 5) aryl,
  - 6) aryl-C<sub>1-10</sub>alkyl,
  - 7) heteroaryl,
  - 8) heteroaryl-C<sub>1-10</sub>alkyl,

20 wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from R<sup>a</sup>; and aryl and heteroaryl optionally substituted with one to four substituents independently selected from R<sup>b</sup>; or

25 R<sup>3</sup>, R<sup>4</sup> and the carbon to which they are attached form a 3-7 membered mono- or bicyclic ring containing 0-2 heteroatoms selected from N, O and S;

- R<sup>5</sup> is
- 1) hydrogen,
  - 30 2) C<sub>1-10</sub> alkyl,
  - 3) Cy, or
  - 4) Cy-C<sub>1-10</sub> alkyl,

wherein alkyl is optionally substituted with one to four substituents independently selected from R<sup>a</sup>; and Cy is optionally substituted with one to four substituents independently selected from R<sup>b</sup>;

- 5 R<sup>6</sup> is           1) hydrogen,  
                  2) C<sub>1-10</sub>alkyl,  
                  3) C<sub>2-10</sub>alkenyl,  
                  4) C<sub>2-10</sub>alkynyl,  
                  5) Cy,  
10               6) Cy-C<sub>1-10</sub>alkyl,  
                  7) Cy-C<sub>2-10</sub>alkenyl,  
                  8) Cy-C<sub>2-10</sub>alkynyl,

wherein alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents selected from R<sup>x</sup>, and Cy is optionally substituted with  
15 one to four substituents independently selected from R<sup>y</sup>;

- R<sup>7</sup> is           1) hydrogen,  
                  2) C<sub>1-10</sub>alkyl,  
                  3) C<sub>2-10</sub>alkenyl,  
20               4) C<sub>2-10</sub>alkynyl,  
                  5) aryl,  
                  6) aryl-C<sub>1-10</sub>alkyl,  
                  7) heteroaryl,  
                  8) heteroaryl-C<sub>1-10</sub>alkyl,

25 wherein alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents selected from R<sub>x</sub>, and aryl and heteroaryl are optionally substituted with one to four substituents independently selected from R<sub>y</sub>; or

30 R<sup>6</sup>, R<sup>7</sup> together with the atoms to which they are attached form a 3-7 membered ring containing 0-2 heteroatoms selected from N, O and S;

R<sup>a</sup> is           1) Cy, or

2) a group selected from Rx;

wherein Cy is optionally substituted with one to four substituents independently selected from R<sup>c</sup>;

- 5 R<sup>b</sup> is
- 1) a group selected from R<sup>a</sup>,
  - 2) C<sub>1-10</sub> alkyl,
  - 3) C<sub>2-10</sub> alkenyl,
  - 4) C<sub>2-10</sub> alkynyl,
  - 5) aryl C<sub>1-10</sub>alkyl,
  - 10 6) heteroaryl C<sub>1-10</sub> alkyl,
- wherein alkyl, alkenyl, alkynyl, aryl, heteroaryl are optionally substituted with a group independently selected from R<sup>c</sup>;

- R<sup>c</sup> is
- 1) halogen,
  - 15 2) amino,
  - 3) carboxy,
  - 4) C<sub>1-4</sub>alkyl,
  - 5) C<sub>1-4</sub>alkoxy,
  - 6) aryl,
  - 20 7) aryl C<sub>1-4</sub>alkyl, or
  - 8) aryloxy;

- R<sup>d</sup> and R<sup>e</sup> are independently selected from hydrogen, C<sub>1-10</sub>alkyl, C<sub>2-10</sub>alkenyl, C<sub>2-10</sub>alkynyl, Cy and Cy C<sub>1-10</sub>alkyl, wherein alkyl, alkenyl, alkynyl and Cy is optionally substituted with one to four substituents independently selected from R<sup>c</sup>; or
- 25 R<sup>d</sup> and R<sup>e</sup> together with the atoms to which they are attached form a heterocyclic ring of 5 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen;

- 30 R<sup>f</sup> and R<sup>g</sup> are independently selected from hydrogen, C<sub>1-10</sub>alkyl, Cy and Cy C<sub>1-10</sub>alkyl; or

$R^f$  and  $R^g$  together with the carbon to which they are attached form a ring of 5 to 7 members containing 0-2 heteroatoms independently selected from oxygen, sulfur and nitrogen;

- 5  $R^h$  is
- 1) hydrogen,
  - 2)  $C_{1-10}$ alkyl,
  - 3)  $C_{2-10}$ alkenyl,
  - 4)  $C_{2-10}$ alkynyl,
  - 5) cyano,
  - 10 6) aryl,
  - 7) aryl  $C_{1-10}$ alkyl,
  - 8) heteroaryl,
  - 9) heteroaryl  $C_{1-10}$ alkyl, or
  - 10)  $-SO_2R^i$ ;
- 15 wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from  $R^a$ ; and aryl and heteroaryl are each optionally substituted with one to four substituents independently selected from  $R^b$ ;
- 20  $R^i$  is
- 1)  $C_{1-10}$ alkyl,
  - 2)  $C_{2-10}$ alkenyl,
  - 3)  $C_{2-10}$ alkynyl, or
  - 4) aryl;
- wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted
- 25 with one to four substituents independently selected from  $R^c$ ;

- $R^x$  is
- 1)  $-OR^d$ ,
  - 2)  $-NO_2$ ,
  - 3) halogen
  - 30 4)  $-S(O)_mR^d$ ,
  - 5)  $-SR^d$ ,
  - 6)  $-S(O)_2OR^d$ ,
  - 7)  $-S(O)_mNR^dR^e$ ,

- 5
- 8)  $-NR^dRe$ ,
  - 9)  $-O(CR^fRg)_nNR^dRe$ ,
  - 10)  $-C(O)R^d$ ,
  - 11)  $-CO_2R^d$ ,
  - 12)  $-CO_2(CR^fRg)_nCONR^dRe$ ,
  - 13)  $-OC(O)R^d$ ,
  - 14)  $-CN$ ,
  - 15)  $-C(O)NR^dRe$ ,
  - 16)  $-NR^dC(O)Re$ ,
  - 10 17)  $-OC(O)NR^dRe$ ,
  - 18)  $-NR^dC(O)ORE$ ,
  - 19)  $-NR^dC(O)NR^dRe$ ,
  - 20)  $-CR^d(N-ORE)$ , or
  - 21)  $-CF_3$ ;

15

- $RY$  is
- 1) a group selected from  $R^x$ ,
  - 2)  $C_{1-10}$  alkyl,
  - 3)  $C_{2-10}$  alkenyl,
  - 4)  $C_{2-10}$  alkynyl,
  - 20 5) aryl  $C_{1-10}$ alkyl,
  - 6) heteroaryl  $C_{1-10}$  alkyl,
  - 7) cycloalkyl,
  - 8) heterocyclyl;

25 wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from  $R^x$ ;  
 Cy is cycloalkyl, heterocyclyl, aryl, or heteroaryl;

$m$  is an integer from 1 to 2;

30  $n$  is an integer from 1 to 10;

- $X$  is
- 1)  $-C(O)OR^d$ ,
  - 2)  $-P(O)_2OR^d$ ,

- 3)  $-S(O)_mOR^d$ ,
- 4)  $-C(O)NR^dR^h$ , or
- 5)  $-5\text{-tetrazolyl}$ .

In one embodiment of the present method compounds of  
 5 formula I are those wherein  $R^1$  is Cy. For the purpose of  $R^1$  Cy is preferably aryl optionally substituted with one to four substituents selected from  $R^b$ . More preferred  $R^1$  is phenyl with a substituent on the 3-position and optionally a second substituent; the more preferred substituents are selected from  $C_{1-10}$ alkoxy, halogen, cyano, and  
 10 trifluoromethyl.

In another embodiment of the present method compounds of Formula I are those wherein  $R^2$  is H or  $C_{1-6}$ alkyl. Preferred  $R^2$  is H or  $C_{1-3}$ alkyl, more preferably H or methyl.

In another embodiment of the present method compounds  
 15 of Formula I are those wherein one of  $R^3$  and  $R^4$  is other than hydrogen. In a preferred embodiment one of  $R^3$  and  $R^4$  is hydrogen and the other is Cy,  $Cy-C_{1-10}$ alkyl, or  $C_{1-10}$ alkyl, each of which is optionally substituted as provided under Formula I. For the purpose of  $R^3/R^4$  Cy is preferably aryl or heteroaryl each optionally substituted with one to four  
 20 substituents selected from  $R^b$ ; and alkyl is optionally substituted with one to four substituents independently selected from  $R^a$ . More preferred  $R^3/R^4$  is selected from phenyl;  $Cy-C_{1-3}$ alkyl wherein Cy is thienyl, pyridyl or phenyl optionally substituted with one or two groups selected from phenyl, hydroxy, halogen, cyano, and nitro; and  $C_{1-6}$ alkyl  
 25 optionally substituted with  $CO_2H$  or  $CONH_2$ .

Yet in another embodiment of the present method compounds of Formula I are those wherein one of  $R^6$  and  $R^7$  is other than hydrogen. In a preferred embodiment one of  $R^6$  and  $R^7$  is hydrogen and the other is aryl,  $aryl-C_{1-10}$ alkyl, or  $C_{1-10}$ alkyl, wherein aryl is  
 30 optionally substituted with one to four substituents independently selected from  $R^y$ , and alkyl is optionally substituted with one to four substituents selected from  $R^x$ . More preferred  $R^6 / R^7$  is selected from


aryl-C<sub>1-3</sub>alkyl wherein aryl is phenyl or naphthyl each optionally substituted with one or two groups selected from R<sup>y</sup>.

Yet in another embodiment of the present method compounds of Formula I are those wherein X is -C(O)OR<sup>d</sup>.

5 In a more preferred embodiment, the cell adhesion is mediated by VLA-4, and the compounds are of Formula Ia as described hereinbelow.

In another aspect the present invention provides compounds of formula Ia:

10



Ia

or a pharmaceutically acceptable salt thereof,

15 wherein

- R<sup>2</sup> is           1)     hydrogen, or  
                  2)     C<sub>1-6</sub>alkyl;

one of R<sup>3</sup> and R<sup>4</sup> is hydrogen and the other is

- 1)     aryl,  
20               2)     heteroaryl,  
                  3)     aryl-C<sub>1-10</sub>alkyl,  
                  4)     heteroaryl-C<sub>1-10</sub>alkyl, or  
                  5)     C<sub>1-10</sub>alkyl,

25 wherein aryl and heteroaryl are each optionally substituted with one to four substituents selected from R<sup>b</sup>; and alkyl is optionally substituted with one to four substituents independently selected from R<sup>a</sup>; or R<sup>3</sup>, R<sup>4</sup> and the carbon to which they are attached form a 3-7 membered mono- or bicyclic ring containing 0-2 heteroatoms selected from N, O and S;


one of R<sup>6</sup> and R<sup>7</sup> is hydrogen and the other is

- 1) aryl,
- 2) aryl-C<sub>1-10</sub>alkyl, or
- 3) C<sub>1-10</sub>alkyl,

- 5 wherein aryl is optionally substituted with one to four substituents independently selected from R<sup>Y</sup>, and alkyl is optionally substituted with one to four substituents selected from R<sup>X</sup>; and R<sup>a</sup>, R<sup>b</sup>, R<sup>x</sup> and R<sup>y</sup> are as defined above under Formula I.


- In a more preferred embodiment of compounds of Formula  
 10 Ia, R<sup>b</sup> is selected from C<sub>1-10</sub>alkoxy, halogen, cyano, and trifluoromethyl; R<sup>2</sup> is H or methyl; one of R<sup>3</sup> and R<sup>4</sup> is hydrogen and the other is phenyl, Cy-C<sub>1-3</sub>alkyl (wherein Cy is thienyl, pyridyl or phenyl optionally substituted with one or two groups selected from phenyl, hydroxy, halogen, cyano, and nitro); and C<sub>1-6</sub>alkyl optionally substituted with  
 15 CO<sub>2</sub>H or CONH<sub>2</sub>; one of R<sup>6</sup> and R<sup>7</sup> is hydrogen and the other is aryl-C<sub>1-2</sub>alkyl, wherein aryl is optionally substituted with one to two substituents independently selected from halogen; and R<sup>y</sup> is as defined above under Formula I.

- The present compounds are generally composed of three  
 20 domains: 1) a sulfonyl moiety, 2) amino acid 1, and 3) amino acid 2, and are named in a manner similar to that used to name oligopeptides. Representative names used herein and their corresponding structures are shown below (without the stereochemistry) to illustrate the nomenclature used in the application.  
 25 N-(3,4-dimethoxybenzenesulfonyl)-(L)-3-(2-thienyl)alanyl-(L)-norleucine




N-(3,5-dichlorobenzenesulfonyl)-(L)-N-methylvalyl-(L)-2-naphthylalanine





N-(3-chlorobenzenesulfonyl)-2-amino-indan-2-carbonyl-(L)-4-fluorophenylalanine

5



"Alkyl", as well as other groups having the prefix "alk", such as alkoxy, alkanoyl, means carbon chains which may be linear or branched or combinations thereof. Examples of alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, sec- and tert-butyl, pentyl, hexyl, heptyl, octyl, nonyl, and the like.

"Alkenyl" means carbon chains which contain at least one carbon-carbon double bond, and which may be linear or branched or combinations thereof. Examples of alkenyl include vinyl, allyl, isopropenyl, pentenyl, hexenyl, heptenyl, 1-propenyl, 2-butenyl, 2-methyl-2-butenyl, and the like.

"Alkynyl" means carbon chains which contain at least one carbon-carbon triple bond, and which may be linear or branched or combinations thereof. Examples of alkynyl include ethynyl, propargyl, 3-methyl-1-pentynyl, 2-heptynyl and the like.

"Cycloalkyl" means mono- or bicyclic saturated carbocyclic rings, each of which having from 3 to 10 carbon atoms. The term also includes monocyclic rings fused to an aryl group in which the point of attachment is on the non-aromatic portion. Examples of cycloalkyl  
5 include cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl, tetrahydronaphthyl, decahydronaphthyl, indanyl, and the like.

"Aryl" means mono- or bicyclic aromatic rings containing only carbon atoms. The term also includes aryl group fused to a monocyclic cycloalkyl or monocyclic heterocyclyl group in which the  
10 point of attachment is on the aromatic portion. Examples of aryl include phenyl, naphthyl, indanyl, indenyl, tetrahydronaphthyl, 2,3-dihydrobenzofuranyl, benzopyranyl, 1,4-benzodioxanyl, and the like.

"Heteroaryl" means a mono- or bicyclic aromatic ring containing at least one heteroatom selected from N, O and S, with each  
15 ring containing 5 to 6 atoms. Examples of heteroaryl include pyrrolyl, isoxazolyl, isothiazolyl, pyrazolyl, pyridyl, oxazolyl, oxadiazolyl, thiadiazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, furanyl, triazinyl, thienyl, pyrimidyl, pyridazinyl, pyrazinyl, benzoxazolyl, benzothiazolyl, benzimidazolyl, benzofuranyl, benzothiophenyl, furo(2,3-b)pyridyl,  
20 quinolyl, indolyl, isoquinolyl, and the like.

"Heterocyclyl" means mono- or bicyclic saturated rings containing at least one heteroatom selected from N, S and O, each of said ring having from 3 to 10 atoms in which the point of attachment may be carbon or nitrogen. The term also includes monocyclic heterocycle fused  
25 to an aryl or heteroaryl group in which the point of attachment is on the non-aromatic portion. Examples of "heterocyclyl" include pyrrolidinyl, piperidinyl, piperazinyl, imidazolidinyl, 2,3-dihydrofuro(2,3-b)pyridyl, benzoxazinyl, tetrahydrohydroquinolinyl, tetrahydroisoquinolinyl, dihydroindolyl, and the like. The term also includes partially  
30 unsaturated monocyclic rings that are not aromatic, such as 2- or 4-pyridones attached through the nitrogen or N-substituted-(1H,3H)-pyrimidine-2,4-diones (N-substituted uracils).

"Halogen" includes fluorine, chlorine, bromine and iodine.

Optical Isomers - Diastereomers - Geometric Isomers - Tautomers

Compounds of Formula I contain one or more asymmetric centers and can thus occur as racemates and racemic mixtures, single  
5 enantiomers, diastereomeric mixtures and individual diastereomers. The present invention is meant to comprehend all such isomeric forms of the compounds of Formula I.

Some of the compounds described herein contain olefinic double bonds, and unless specified otherwise, are meant to include both  
10 E and Z geometric isomers.

Some of the compounds described herein may exist with different points of attachment of hydrogen, referred to as tautomers. Such an example may be a ketone and its enol form known as keto-enol  
15 tautomers. The individual tautomers as well as mixture thereof are encompassed with compounds of Formula I.

Compounds of the Formula I may be separated into diastereoisomeric pairs of enantiomers by, for example, fractional crystallization from a suitable solvent, for example methanol or ethyl acetate or a mixture thereof. The pair of enantiomers thus obtained may  
20 be separated into individual stereoisomers by conventional means, for example by the use of an optically active acid as a resolving agent.

Alternatively, any enantiomer of a compound of the general Formula I or Ia may be obtained by stereospecific synthesis using optically pure starting materials or reagents of known configuration.

25

Salts

The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids including inorganic or organic bases and inorganic or organic acids.  
30 Salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic salts, manganous, potassium, sodium, zinc, and the like. Particularly preferred are the ammonium, calcium, magnesium, potassium, and

sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic base include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, and basic ion exchange resins, such as arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethyl-morpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine, and the like.

When the compound of the present invention is basic, salts may be prepared from pharmaceutically acceptable non-toxic acids, including inorganic and organic acids. Such acids include acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic acid, and the like. Particularly preferred are citric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric, and tartaric acids.

It will be understood that, as used herein, references to the compounds of Formula I are meant to also include the pharmaceutically acceptable salts.

#### Utilities

The ability of the compounds of Formula I to antagonize the actions of VLA-4 and/or  $\alpha 4\beta 7$  integrin makes them useful for preventing or reversing the symptoms, disorders or diseases induced by the binding of VLA-4 and/or  $\alpha 4\beta 7$  to their various respective ligands. Thus, these antagonists will inhibit cell adhesion processes including cell activation, migration, proliferation and differentiation. Accordingly, another aspect of the present invention provides a method for the treatment

(including prevention, alleviation, amelioration or suppression) of diseases or disorders or symptoms mediated by VLA-4 and/or  $\alpha 4\beta 7$  binding and cell adhesion and activation, which comprises administering to a mammal an effective amount of a compound of

5 Formula I. Such diseases, disorders, conditions or symptoms are for example (1) multiple sclerosis, (2) asthma, (3) allergic rhinitis, (4) allergic conjunctivitis, (5) inflammatory lung diseases, (6) rheumatoid arthritis, (7) septic arthritis, (8) type I diabetes, (9) organ transplantation rejection, (10) restenosis, (11) autologous bone marrow transplantation,

10 (12) inflammatory sequelae of viral infections, (13) myocarditis, (14) inflammatory bowel disease including ulcerative colitis and Crohn's disease, (15) certain types of toxic and immune-based nephritis, (16) contact dermal hypersensitivity, (17) psoriasis, (18) tumor metastasis, and (19) atherosclerosis.

15

#### Dose Ranges

The magnitude of prophylactic or therapeutic dose of a compound of Formula I will, of course, vary with the nature of the severity of the condition to be treated and with the particular compound

20 of Formula I and its route of administration. It will also vary according to the age, weight and response of the individual patient. In general, the daily dose range lie within the range of from about 0.001 mg to about 100 mg per kg body weight of a mammal, preferably 0.01 mg to about 50 mg per kg, and most preferably 0.1 to 10 mg per kg, in single or divided

25 doses. On the other hand, it may be necessary to use dosages outside these limits in some cases.

For use where a composition for intravenous administration is employed, a suitable dosage range is from about 0.001 mg to about 25 mg (preferably from 0.01 mg to about 1 mg) of a compound

30 of Formula I per kg of body weight per day and for cytoprotective use from about 0.1 mg to about 100 mg (preferably from about 1 mg to about 100 mg and more preferably from about 1 mg to about 10 mg) of a compound of Formula I per kg of body weight per day.

In the case where an oral composition is employed, a suitable dosage range is, e.g. from about 0.01 mg to about 100 mg of a compound of Formula I per kg of body weight per day, preferably from about 0.1 mg to about 10 mg per kg and for cytoprotective use from 0.1 mg  
5 to about 100 mg (preferably from about 1 mg to about 100 mg and more preferably from about 10 mg to about 100 mg) of a compound of Formula I per kg of body weight per day.

For the treatment of diseases of the eye, ophthalmic preparations for ocular administration comprising 0.001-1% by weight  
10 solutions or suspensions of the compounds of Formula I in an acceptable ophthalmic formulation may be used.

#### Pharmaceutical Compositions

Another aspect of the present invention provides  
15 pharmaceutical compositions which comprises a compound of Formula I and a pharmaceutically acceptable carrier. The term "composition", as in pharmaceutical composition, is intended to encompass a product comprising the active ingredient(s), and the inert ingredient(s) (pharmaceutically acceptable excipients) that make up the carrier, as  
20 well as any product which results, directly or indirectly, from combination, complexation or aggregation of any two or more of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients. Accordingly, the pharmaceutical compositions of the  
25 present invention encompass any composition made by admixing a compound of Formula I, additional active ingredient(s), and pharmaceutically acceptable excipients.

Any suitable route of administration may be employed for providing a mammal, especially a human with an effective dosage of a  
30 compound of the present invention. For example, oral, rectal, topical, parenteral, ocular, pulmonary, nasal, and the like may be employed. Dosage forms include tablets, troches, dispersions, suspensions, solutions, capsules, creams, ointments, aerosols, and the like.

The pharmaceutical compositions of the present invention comprise a compound of Formula I as an active ingredient or a pharmaceutically acceptable salt thereof, and may also contain a pharmaceutically acceptable carrier and optionally other therapeutic ingredients. The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids including inorganic bases or acids and organic bases or acids.

The compositions include compositions suitable for oral, rectal, topical, parenteral (including subcutaneous, intramuscular, and intravenous), ocular (ophthalmic), pulmonary (aerosol inhalation), or nasal administration, although the most suitable route in any given case will depend on the nature and severity of the conditions being treated and on the nature of the active ingredient. They may be conveniently presented in unit dosage form and prepared by any of the methods well-known in the art of pharmacy.

For administration by inhalation, the compounds of the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or nebulisers. The compounds may also be delivered as powders which may be formulated and the powder composition may be inhaled with the aid of an insufflation powder inhaler device. The preferred delivery systems for inhalation are metered dose inhalation (MDI) aerosol, which may be formulated as a suspension or solution of a compound of Formula I in suitable propellants, such as fluorocarbons or hydrocarbons and dry powder inhalation (DPI) aerosol, which may be formulated as a dry powder of a compound of Formula I with or without additional excipients.

Suitable topical formulations of a compound of formula I include transdermal devices, aerosols, creams, ointments, lotions, dusting powders, and the like.

In practical use, the compounds of Formula I can be combined as the active ingredient in intimate admixture with a pharmaceutical carrier according to conventional pharmaceutical

compounding techniques. The carrier may take a wide variety of forms depending on the form of preparation desired for administration, e.g., oral or parenteral (including intravenous). In preparing the compositions for oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like in the case of oral liquid preparations, such as, for example, suspensions, elixirs and solutions; or carriers such as starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants, binders, disintegrating agents and the like in the case of oral solid preparations such as, for example, powders, capsules and tablets, with the solid oral preparations being preferred over the liquid preparations. Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit form in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be coated by standard aqueous or nonaqueous techniques.

In addition to the common dosage forms set out above, the compounds of Formula I may also be administered by controlled release means and/or delivery devices such as those described in U.S. Patent Nos. 3,845,770; 3,916,899; 3,536,809; 3,598,123; 3,630,200 and 4,008,719.

Pharmaceutical compositions of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient, as a powder or granules or as a solution or a suspension in an aqueous liquid, a non-aqueous liquid, an oil-in-water emulsion or a water-in-oil liquid emulsion. Such compositions may be prepared by any of the methods of pharmacy but all methods include the step of bringing into association the active ingredient with the carrier which constitutes one or more necessary ingredients. In general, the compositions are prepared by uniformly and intimately admixing the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product into the desired presentation. For example, a tablet may be prepared by compression or



molding, optionally with one or more accessory ingredients.

Compressed tablets may be prepared by compressing in a suitable machine, the active ingredient in a free-flowing form such as powder or granules, optionally mixed with a binder, lubricant, inert diluent,

- 5 surface active or dispersing agent. Molded tablets may be made by molding in a suitable machine, a mixture of the powdered compound moistened with an inert liquid diluent. Desirably, each tablet contains from about 1 mg to about 500 mg of the active ingredient and each cachet or capsule contains from about 1 to about 500 mg of the active ingredient.

- 10 The following are examples of representative pharmaceutical dosage forms for the compounds of Formula I:

Injectable Suspension (I.M.) mg/mL

	Compound of Formula I	10
15	Methylcellulose	5.0
	Tween 80	0.5
	Benzyl alcohol	9.0
	Benzalkonium chloride	1.0
	Water for injection to a total volume of 1 mL	

20

Tablet mg/tablet

	Compound of Formula I	25
	Microcrystalline Cellulose	415
	Povidone	14.0
25	Pregelatinized Starch	43.5
	Magnesium Stearate	<u>2.5</u>
		500

Capsule mg/capsule

30	Compound of Formula I	25
	Lactose Powder	573.5
	Magnesium Stearate	<u>1.5</u>
		600

	<u>Aerosol</u>	<u>Per canister</u>
	Compound of Formula I	24 mg
	Lecithin, NF Liquid Concentrate	1.2 mg
5	Trichlorofluoromethane, NF	4.025 g
	Dichlorodifluoromethane, NF	12.15 g

#### Combination Therapy

- Compounds of Formula I may be used in combination with other drugs that are used in the treatment/prevention/suppression or amelioration of the diseases or conditions for which compounds of Formula I are useful. Such other drugs may be administered, by a route and in an amount commonly used therefor, contemporaneously or sequentially with a compound of Formula I. When a compound of Formula I is used contemporaneously with one or more other drugs, a pharmaceutical composition containing such other drugs in addition to the compound of Formula I is preferred. Accordingly, the pharmaceutical compositions of the present invention include those that also contain one or more other active ingredients, in addition to a compound of Formula I. Examples of other active ingredients that may be combined with a compound of Formula I, either administered separately or in the same pharmaceutical compositions, include, but are not limited to:
- (a) other VLA-4 antagonists such as those described in US 5,510,332, WO97/03094, WO97/02289, WO96/40781, WO96/22966, WO96/20216, WO96/01644, WO96/06108, WO95/15973 and WO96/31206;
  - (b) steroids such as beclomethasone, methylprednisolone, betamethasone, prednisone, dexamethasone, and hydrocortisone;
  - (c) immunosuppressants such as cyclosporin, tacrolimus, rapamycin and other FK-506 type immunosuppressants;
  - (d) antihistamines (H1-histamine antagonists) such as bromopheniramine, chlorpheniramine, dexchlorpheniramine, triprolidine, clemastine, diphenhydramine, diphenylpyraline, tripeleminamine, hydroxyzine, methdilazine, promethazine,


- trimeprazine, azatadine, cyproheptadine, antazoline, pheniramine  
pyrilamine, astemizole, terfenadine, loratadine, cetirizine,  
fexofenadine, descarboethoxyloratadine, and the like; (e) non-steroidal  
anti-asthmatics such as  $\beta$ 2-agonists (terbutaline, metaproterenol,  
5 fenoterol, isoetharine, albuterol, bitolterol, and pirbuterol), theophylline,  
cromolyn sodium, atropine, ipratropium bromide, leukotriene  
antagonists (zafirlukast, montelukast, pranlukast, iralukast,  
pobilukast, SKB-106,203), leukotriene biosynthesis inhibitors (zileuton,  
BAY-1005); (f) non-steroidal antiinflammatory agents (NSAIDs) such as  
10 propionic acid derivatives (alminoprofen, benoxaprofen, bucloxic acid,  
carprofen, fenbufen, fenoprofen, fluprofen, flurbiprofen, ibuprofen,  
indoprofen, ketoprofen, miroprofen, naproxen, oxaprozin, pirprofen,  
pranoprofen, suprofen, tiaprofenic acid, and tioxaprofen), acetic acid  
derivatives (indomethacin, acetaminophen, alclofenac, clidanac, diclofenac,  
15 fenclofenac, fenclozic acid, fentiazac, furofenac, ibufenac, isoxepac,  
oxipinac, sulindac, tiopinac, tolmetin, zidometacin, and zomepirac),  
fenamic acid derivatives (flufenamic acid, meclofenamic acid,  
mefenamic acid, niflumic acid and tolfenamic acid), biphenylcarboxylic  
acid derivatives (diflunisal and flufenisal), oxicams (isoxicam,  
20 piroxicam, sudoxicam and tenoxicam), salicylates (acetyl salicylic acid,  
sulfasalazine) and the pyrazolones (apazone, bezpiperylon, feprazone,  
mofebutazone, oxyphenbutazone, phenylbutazone); (g) cyclooxygenase-2  
(COX-2) inhibitors such as celecoxib; (h) inhibitors of phosphodiesterase  
type IV (PDE-IV); (i) antagonists of the chemokine receptors, especially  
25 CCR-1, CCR-2, and CCR-3; (j) cholesterol lowering agents such as HMG-  
CoA reductase inhibitors (lovastatin, simvastatin and pravastatin,  
fluvastatin, atorvastatin, and other statins), sequestrants  
(cholestyramine and colestipol), nicotinic acid, fenofibric acid derivatives  
(gemfibrozil, clofibrat, fenofibrate and benazafibrate), and probucol; (k)  
30 anti-diabetic agents such as insulin, sulfonylureas, biguanides  
(metformin),  $\alpha$ -glucosidase inhibitors (acarbose) and glitazones  
(troglitazone, pioglitazone, englitazone, MCC-555, BRL49653 and the  
like); (l) preparations of interferon beta (interferon beta-1a, interferon

beta-1b); (m) anticholinergic agents such as muscarinic antagonists (ipratropium bromide); (n) other compounds such as 5-aminosalicylic acid and prodrugs thereof, antimetabolites such as azathioprine and 6-mercaptapurine, and cytotoxic cancer chemotherapeutic agents.

5           The weight ratio of the compound of the Formula I to the second active ingredient may be varied and will depend upon the effective dose of each ingredient. Generally, an effective dose of each will be used. Thus, for example, when a compound of the Formula I is combined with an NSAID the weight ratio of the compound of the  
10   Formula I to the NSAID will generally range from about 1000:1 to about 1:1000, preferably about 200:1 to about 1:200. Combinations of a compound of the Formula I and other active ingredients will generally also be within the aforementioned range, but in each case, an effective dose of each active ingredient should be used.

15           Compounds of the present invention may be prepared by procedures illustrated in the accompanying schemes. In the first method (Scheme 1), a resin-based synthetic strategy is outlined where the resin employed is represented by the ball (●). An N-Fmoc-protected amino acid derivative **A** (Fmoc = fluorenylmethoxycarbonyl) is loaded on  
20   to the appropriate hydroxyl-containing resin using dicyclohexylcarbodiimide (DCC) and 1-hydroxybenzotriazole (HOBt) in dimethylformamide (DMF) to give **B**. The Fmoc protecting group is removed with piperidine in DMF to yield free amine **C**. The next Fmoc-protected amino acid derivative **D** is coupled to **C** employing standard  
25   peptide (in this instance, 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU), HOBt, and N,N-diisopropylethylamine (DIEA) in DMF) to yield dipeptide **E**. The Fmoc group is removed with piperidine in DMF to yield the free amine **F**. A sulfonyl chloride derivative is reacted with **F** in the presence of DIEA to  
30   yield **G**. The final product is removed from the resin with strong acid (in this instance, trifluoroacetic acid (TFA) in the presence of thioanisole and dithiane) to yield compounds of the present invention **H**.


Scheme 1



In the second method (Scheme 2), standard solution phase synthetic methodology is outlined. An N-Boc-protected amino acid derivative **A** (Boc = tert-butyloxycarbonyl) is treated with tert-butyl 2,2,2-trichloroacetimidate in the presence of boron trifluoride etherate to yield tert-butyl ester **B** which is subsequently coupled to Cbz-protected amino acid derivative **C** (Cbz = carbobenzyloxy) in the presence of 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC), HOBT, and N-methylmorpholine (NMM) in methylene chloride (CH<sub>2</sub>Cl<sub>2</sub>) to yield dipeptide **D**. Catalytic hydrogenation of **D** in the presence of a palladium-on-carbon (Pd/C) catalyst yields **E**. Reaction of **E** with a sulfonyl chloride in the presence of DIEA and 4-dimethylaminopyridine (DMAP) yields **F** which is subsequently reacted with strong acid (TFA) to yield the desired product **G**.

15

Scheme 2



GENERAL PROCEDURE FOR THE SOLID-PHASE SYNTHESIS OF  
COMPOUNDS OF FORMULA I.

Step A. Loading of N-Fmoc-amino acid derivatives onto resins.

- 5           N-Fmoc-amino acids were loaded on either Wang®  
(Calbiochem-Novabiochem Corp.) or Chloro (2-chlorotriyl) resin.  
Wang® resin, typically 0.3 mmol, was washed with dimethylformamide  
three times. A solution of N-Fmoc-amino acid (0.3 mmol) in  
dimethylformamide (3 mL) was transferred to the pre-swollen Wang®  
10 resin. Dicyclohexylcarbodiimide (0.3 mmol) and 1-N-  
hydroxybenztriazole (0.3 mmol) was added and the mixture gently  
swirled for 2 hours. Following filtration, the resin was sequentially  
washed with dimethylformamide (3 times) and dichloromethane (3  
times). The amino acid substitution value obtained after vacuum drying  
15 typically ranged between 0.07 to 0.1 mmol.

- Alternatively, Chloro (2-chlorotriyl) resin, typically 0.2  
mmol, was pre-swollen in dimethylformamide. A solution of N-Fmoc-  
amino acid (0.2 mmol) in dimethylformamide (3 ml) was added to the  
resin, followed by the addition of N,N-diisopropylethylamine(0.4 mmol).  
20 The resin was gently stirred for 2 hours, filtered and washed  
sequentially with dimethylformamide (3 times) and dichloromethane (3  
times). The resin was finally washed with 10% methanol in  
dichloromethane and vacuum dried. The amino acid substitution value  
obtained after vacuum drying typically ranged between 0.05 to 0.1 mmol.

25

Step B. Deprotection of the N-Fmoc group.

- The N-Fmoc protecting group was removed from the resin  
from Step A by treatment with 20% piperidine in dimethylformamide for  
30 minutes. Following filtration, the resin was washed sequentially  
30 with dimethylformamide (3 times), dichloromethane (1 time) and  
dimethylformamide (2 times) and used in the subsequent reaction.



Step C. Coupling of the next N-Fmoc-amino acid derivative

A solution of the next desired N-Fmoc-amino acid derivative (0.4 mmol) in dimethylformamide (2 mL) was mixed with 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (0.4 mmol), 1-N-hydroxybenzotriazole (0.4 mmol) and diisopropylethylamine (0.6 mmol). This solution was transferred to resin from Step B and typically allowed to react for 2 hours. Couplings were monitored by ninhydrin reaction. The coupling mixture was filtered and the resin washed with dimethylformamide (3 times) and used in the subsequent reaction.

Step D. Deprotection of the N-Fmoc group.

The N-Fmoc protecting group was removed from the resin from Step C by the procedure described in Step B and used in the subsequent reaction.

Step E. Acylation (or sulfonylation) of the terminal amino group.

The desired N-terminal capping reagent (sulfonylchloride or acylchloride) (0.4 mol) was dissolved in dimethylformamide (2 ml), mixed with N,N-diisopropylethylamine (0.8 mmol) and added to the resin from Step D. After approximately two hours, the resin was sequentially washed with dimethylformamide (3 times) and dichloromethane (3 times).

Step F. Cleavage of the desired products from the resins.

The final desired products were cleaved from the resins from Step E by gently stirring with a solution of trifluoroacetic acid:thioanisole:ethanedithiol (95:2.5:2.5); 3 hours for Wang® resin and 30 minutes for the Chloro (2-chlorotriyl) resin. Following filtration, the solvents were removed by evaporation and the residue dissolved in acetonitrile (3 mL). Insoluble material was removed by filtration. The final products were purified by reverse phase chromatography with a linear gradient of buffer A (0.1% trifluoroacetic acid in water) and buffer

B (0.1% trifluoroacetic acid in acetonitrile) and isolated by lyophilization. Molecular ions were obtained by electrospray ionization mass spectrometry or matrix-assisted laser desorption ionization time-of-flight mass spectrometry to confirm the structure of each peptide.

- 5                   The following compounds were prepared by the general procedure described above using the appropriate amino acid derivatives and sulfonyl chloride. These examples are provided to illustrate the present invention and are not to be construed as limiting its scope in any manner.

10

<u>Ex. No.</u>	<u>Compound Name</u>	<u>MS*</u>
(1)	N-(3,4-dimethoxybenzenesulfonyl)-(L)-3-(2-thienyl)alanyl-norleucine;	485
(2)	N-(3,4-dimethoxybenzenesulfonyl)-(L)-3,3-diphenylalanyl-(L)-norleucine;	555
(3)	N-(3,4-dimethoxybenzenesulfonyl)-(L)-aspartyl-(L)-norleucine;	447
(4)	N-(3,4-dimethoxybenzenesulfonyl)-(L)-3-(3-pyridyl)alanyl-(L)-norleucine;	481
(5)	N-(3,4-dimethoxybenzenesulfonyl)-(L)-alanyl-(L)-norleucine;	403
(6)	N-(3,4-dimethoxybenzenesulfonyl)-(L)-homophenylalanyl-(L)-norleucine;	493
(7)	N-(3,4-dimethoxybenzenesulfonyl)-(L)-valyl-(L)-3-(2-naphthyl)alanine;	515
(8)	N-(3,4-dimethoxybenzenesulfonyl)-(L)-asparagyl-(L)-norleucine;	446
(9)	N-(3,4-dimethoxybenzenesulfonyl)-(L)-isoleucyl-(L)-3-(2-naphthyl)alanine;	529
(10)	N-(3,4-dimethoxybenzenesulfonyl)-(L)-3-(4-biphenyl)alanyl-(L)-norleucine;	555

(11)	N-(3,5-dichlorobenzenesulfonyl)-(L)-N-methylvalyl-(L)-3-(2-naphthyl)alanine;	537
(12)	N-(3,5-di(trifluoromethyl)-benzenesulfonyl)-(L)-N-methylvalyl-(L)-3-(2-naphthyl)alanine;	605
(13)	N-(3,5-dichlorobenzenesulfonyl)-2(S)-aminobutyryl-(L)-3-(2-naphthyl)alanine;	509
(14)	N-(3,4-dimethoxybenzenesulfonyl)-(L)-N-methylisoleucyl-(L)-3-(2-naphthyl)alanine;	543
(15)	N-(3,4-dimethoxybenzenesulfonyl)-(L)-N-methylvalyl-(L)-3-(2-naphthyl)alanine;	529
(16)	N-(3,4-dimethoxybenzenesulfonyl)-(L)-sarcosinyl-(L)-3-(2-naphthyl)alanine;	487
(17)	N-(3,4-dimethoxybenzenesulfonyl)-(L)-phenylalanyl-(L)-norleucine;	479
(18)	N-(3,4-dimethoxybenzenesulfonyl)-(L)-N-methylphenylalanyl-(L)-norleucine;	492
(19)	N-(3,4-dimethoxybenzenesulfonyl)-(L)- $\alpha$ -phenylglycyl-(L)-norleucine;	465
(20)	N-(3,4-dimethoxybenzenesulphonyl)-(L)-N-methylalanyl-(L)-norleucine;	417
(21)	N-(3,5-dichlorobenzenesulphonyl)-(L)-tyrosyl-(L)-4-fluorophenylalanine;	555
(22)	N-(3,5-dichlorobenzenesulphonyl)-(L)-3-(4-pyridyl)alanine-(L)-4-fluorophenylalanine;	540
(23)	N-(3,5-dichlorobenzenesulphonyl)-(L)-phenylalanyl-(L)-4-fluorophenylalanine;	539.1
(24)	N-(3,5-dichlorobenzenesulphonyl)-(L)-4-fluorophenylalanyl-(L)-4-fluorophenylalanine;	557.1
(25)	N-(3,5-dichlorobenzenesulphonyl)-(L)-3,5-difluorophenylalanyl-(L)-4-fluorophenylalanine;	575.1
(26)	N-(3,5-dichlorobenzenesulphonyl)-(L)-3-(2-pyridyl)alanyl-(L)-4-fluorophenylalanine;	540
(27)	N-(3,5-dichlorobenzenesulphonyl)-(L)-N-	543.3

	methylasspartyl-(L)-3-(2-naphthyl)alanine;	
(28)	N-(3-fluorobenzenesulphonyl)-(L)-N-methylvalyl-(L)-3-(2-naphthyl)alanine;	487.2
(29)	N-(3,5-dichlorobenzenesulphonyl)-(L)-4-chlorophenylalanyl-(L)-4-fluorophenylalanine;	575.1
(30)	N-(3,5-dichlorobenzenesulphonyl)-(L)-4-cyanophenylalanyl-(L)-4-fluorophenylalanine;	564.0
(31)	N-(3,5-dichlorobenzenesulphonyl)-(L)-4-nitrophenylalanyl-(L)-4-fluorophenylalanine;	584.1
(32)	N-(3,5-dichlorobenzenesulphonyl)-(L)-aspartyl-(L)-4-fluorophenylalanine;	507.0
(33)	N-(3,5-dichlorobenzenesulphonyl)-(L)-3-(3-pyridyl)alanyl-(L)-4-fluorophenylalanine;	540.2
(34)	N-(3,5-dichlorobenzenesulphonyl)-(L)-N-methylvalyl-(L)-4-fluorophenylalanine;	604.2
(35)	N-(3,5-dichlorobenzenesulphonyl)-(D)-N-methylvalyl-(L)-3-(2-naphthyl)alanine;	537.0
(36)	N-(3-chlorobenzenesulphonyl)-2-amino-indan-2-carbonyl-(L)-4-fluorophenylalanine;	517.2
(37)	N-(3-fluorobenzenesulphonyl)-(L)-2-phenylglycyl-(L)-4-fluorophenylalanine;	475.2
(38)	N-(3,5-dichlorobenzenesulphonyl)-(L)-leucyl-(L)-aspartic acid.	455.1

\*m/e, (M)<sup>+</sup> or (M + 1)<sup>+</sup> or (M + NH<sub>4</sub>)<sup>+</sup>.

#### EXAMPLE 39

##### 5 Inhibition of VLA-4 Dependent Adhesion to BSA-CS-1 Conjugate

###### Step 1. Preparation of CS-1 Coated Plates

10 Untreated 96 well polystyrene flat bottom plates were coated with bovine serum albumin (BSA; 20 µg/ml) for 2 hours at room

temperature and washed twice with phosphate buffered saline (PBS). The albumin coating was next derivatized with 10 µg/ml 3-(2-pyridyldithio) propionic acid N-hydroxysuccinimide ester (SPDP), a heterobifunctional crosslinker, for 30 minutes at room temperature and washed twice with PBS. The CS-1 peptide (Cys-Leu-His-Gly-Pro-Glu-Ile-Leu-Asp-Val-Pro-Ser-Thr), which was synthesized by conventional solid phase chemistry and purified by reverse phase HPLC, was next added to the derivatized BSA at a concentration of 2.5 µg/ml and allowed to react for 2 hours at room temperature. The plates were washed twice with PBS and stored at 4°C.

#### Step 2. Preparation of Fluorescently Labeled Jurkat Cells

Jurkat cells, clone E6-1, obtained from the American Type Culture Collection (Rockville, MD; cat # ATCC TIB-152) were grown and maintained in RPMI-1640 culture medium containing 10% fetal calf serum (FCS), 50 units/ml penicillin, 50 µg/ml streptomycin and 2 mM glutamine. Fluorescence activated cell sorter analysis with specific monoclonal antibodies confirmed that the cells expressed both the α4 and β1 chains of VLA-4. The cells were centrifuged at 400xg for five minutes and washed twice with PBS. The cells were incubated at a concentration of  $2 \times 10^6$  cells/ml in PBS containing a 1 µM concentration of a fluorogenic esterase substrate (2', 7'-bis-(2-carboxyethyl)-5-(and -6)-carboxyfluorescein, acetoxymethyl ester; BCECF-AM; Molecular Probes Inc., Eugene, Oregon; catalog #B-1150) for 30-60 minutes at 37°C in a 5% CO<sub>2</sub>/air incubator. The fluorescently labeled Jurkat cells were washed two times in PBS and resuspended in RPMI containing 0.25% BSA at a final concentration of  $2.0 \times 10^6$  cells/ml.

#### Step 3. Assay Procedure

Compounds of this invention were prepared in DMSO at 100x the desired final assay concentration. Final concentrations were

selected from a range between 0.001 nM-100  $\mu$ M. Three  $\mu$ L of diluted compound, or vehicle alone, were premixed with 300  $\mu$ L of cell suspension in 96-well polystyrene plates with round bottom wells. 100  $\mu$ L aliquots of the cell /compound mixture were then transferred in duplicate to CS-1 coated wells. The cells were next incubated for 30 minutes at room temperature. The non-adherent cells were removed by two gentle washings with PBS. The remaining adherent cells were quantitated by reading the plates on a Cytofluor II fluorescence plate reader (Perseptive Biosystems Inc., Framingham, MA; excitation and emission filter settings were 485 nm and 530 nm, respectively). Control wells containing vehicle alone were used to determine the level of cell adhesion corresponding to 0% inhibition. Control wells coated with BSA and crosslinker (no CS-1 peptide) were used to determine the level of cell adhesion corresponding to 100% inhibition. Cell adhesion to wells coated with BSA and crosslinker was usually less than 5% of that observed to CS-1 coated wells in the presence of vehicle. Percent inhibition was then calculated for each test well and the  $IC_{50}$  was determined from a ten point titration using a validated four parameter fit algorithm.

20

## EXAMPLE 40

Antagonism of VLA-4 Dependent Binding to VCAM-Ig Fusion Protein.

## 1. Preparation of VCAM-Ig

25

The signal peptide as well as domains 1 and 2 of human VCAM (GenBank Accession no. M30257) were amplified by PCR using the human VCAM cDNA (R & D Systems) as template and the following primer sequences: 3'-PCR primer: 5'-AATTATAATTGATCAACTTAC CTGTCAATTCTTTTACAGCCTGCC-3';

30

5'-PCR primer:

5'-ATAGGAATTCCAGCTGCCACCATGCCTGGGAAGATGGTTCG-3'.

The 5'-PCR primer contained EcoRI and PvuII restriction sites followed by a Kozak consensus sequence (CCACC) proximal to the

initiator methionine ATG. The 3'-PCR primer contained a BclI site and a splice donor sequence. PCR was performed for 30 cycles using the following parameters: 1 min. at 94°C, 2 min. at 55°C, and 2 min. at 72°C. The amplified region encoded the following sequence of human

- 5 VCAM-1:  
MPGKMVVILGASNILWIMFAASQAFKIETTPESRYLAQIGDSVSLTC  
STTGCESPFFFSWRTQIDSPLNGKVTNEGTTSTLTMPVSVFGNEHSYLC  
TATCESRKLEKGIQVEIYSFPKDP EIHLSGPLEAGKPITVKCSVADVY  
PFDRLEIDLLKGDHLMKSQEFLEDADRKSLETKSLEVTFTPVIEDIGKV  
10 LVCRAKLHIDEMDSVPTVRQAVKEL. The resulting PCR product of  
650 bp was digested with EcoRI and BclI and ligated to expression vector  
pIg-Tail (R & D Systems, Minneapolis, MN) digested with EcoRI and  
BamHI. The pIg-Tail vector contains the genomic fragment which  
encodes the hinge region, CH2 and CH3 of human IgG1 (GenBank  
15 Accession no. Z17370). The DNA sequence of the resulting VCAM  
fragment was verified using Sequenase (US Biochemical, Cleveland,  
OH). The fragment encoding the entire VCAM-Ig fusion was  
subsequently excised from pIg-Tail with EcoRI and NotI and ligated to  
pCI-neo (Promega, Madison, WI) digested with EcoRI and NotI. The  
20 resulting vector, designated pCI-neo/VCAM-Ig was transfected into  
CHO-K1 (ATCC CCL 61) cells using calcium-phosphate DNA  
precipitation (Specialty Media, Lavallete, NJ). Stable VCAM-Ig  
producing clones were selected according to standard protocols using  
0.2-0.8 mg/ml active G418 (Gibco, Grand Island, NY), expanded, and cell  
25 supernatants were screened for their ability to mediate Jurkat adhesion  
to wells previously coated with 1.5 µg/ml (total protein) goat anti-human  
IgG (Sigma, St. Louis, MO). A positive CHO-K1/VCAM-Ig clone was  
subsequently adapted to CHO-SFM serum-free media (Gibco) and  
maintained under selection for stable expression of VCAM-Ig. VCAM-  
30 Ig was purified from crude culture supernatants by affinity  
chromatography on Protein A/G Sepharose (Pierce, Rockford, IL)  
according to the manufacturer's instructions and desalted into 50 mM

sodium phosphate buffer, pH 7.6, by ultrafiltration on a YM-30 membrane (ALicon, Beverly, MA).

Step 2. Preparation of  $^{125}\text{I}$ -VCAM-Ig

- 5 VCAM-Ig was labeled to a specific radioactivity greater than 1000 Ci/mmol with  $^{125}\text{I}$ -Bolton Hunter reagent (New England Nuclear, Boston, MA; cat # NEX120-0142) according to the manufacturer's instructions. The labeled protein was separated from unincorporated isotope by means of a calibrated HPLC gel filtration column (G2000SW; 10 7.5 x 600 mm; Tosoh, Japan) using uv and radiometric detection.

Step 3. VCAM-Ig Binding Assay

- Compounds of this invention were prepared in DMSO at 100x the desired final assay concentration. Final concentrations were 15 selected from a range between 0.001 nM-100  $\mu\text{M}$ . Jurkat cells were centrifuged at 400xg for five minutes and resuspended in binding buffer (25 mM HEPES, 150 mM NaCl, 3 mM KCl, 2 mM glucose, 0.1% bovine serum albumin, pH 7.4). The cells were centrifuged again and resuspended in binding buffer supplemented with  $\text{MnCl}_2$  at a final 20 concentration of 1 mM. Compounds were assayed in Millipore MHVB multiscreen plates (cat# MHVBN4550, Millipore Corp., MA) by making the following additions to duplicate wells: (i) 200  $\mu\text{L}$  of binding buffer containing 1 mM  $\text{MnCl}_2$ ; (ii) 20  $\mu\text{L}$  of  $^{125}\text{I}$ -VCAM-Ig in binding buffer containing 1 mM  $\text{MnCl}_2$  (final assay concentration ~ 100 pM); (iii) 2.5  $\mu\text{L}$  25 of compound solution or DMSO; (iv) and  $0.5 \times 10^6$  cells in a volume of 30  $\mu\text{L}$ . The plates were incubated at room temperature for 30 minutes, filtered on a vacuum box, and washed on the same apparatus by the addition of 100  $\mu\text{L}$  of binding buffer containing 1 mM  $\text{MnCl}_2$ . After insertion of the multiscreen plates into adapter plates (Packard, 30 Meriden, CT, cat# 6005178), 100  $\mu\text{L}$  of Microscint-20 (Packard cat# 6013621) was added to each well. The plates were then sealed, placed on a shaker for 30 seconds, and counted on a Topcount microplate



scintillation counter (Packard). Control wells containing DMSO alone were used to determine the level of VCAM-Ig binding corresponding to 0% inhibition. Control wells in which cells were omitted were used to determine the level of binding corresponding to 100% inhibition. Binding of  $^{125}\text{I}$ -VCAM-Ig in the absence of cells was usually less than 5% of that observed using cells in the presence of vehicle. Percent inhibition was then calculated for each test well and the  $\text{IC}_{50}$  was determined from a ten point titration using a validated four parameter fit algorithm.

10

## EXAMPLE 41

Antagonism of  $\alpha_4\beta_1$  Dependent Binding to VCAM-Ig Fusion Protein.

Step 1.  $\alpha_4\beta_1$  Cell line.

15

RPMI-8866 cells (a human B cell line  $\alpha_4\beta_1\beta_2^+$ ; a gift from Prof. John Wilkins, University of Manitoba, Canada) were grown in RPMI/10% fetal calf serum/ 100 U penicillin/100  $\mu\text{g}$  streptomycin/2 mM L-glutamine at  $37^\circ\text{C}$ , 5 % carbon dioxide. The cells were pelleted at 1000 rpm for 5 minutes and then washed twice and resuspended in binding buffer (25 mM Hepes, 150 mM NaCl, 0.1 % BSA, 3 mM KCl, 2 mM Glucose, pH 7.4).

20

Step 2. VCAM-Ig Binding Assay

25

Compounds of this invention were prepared in DMSO at 100x the desired final assay concentration. Final concentrations were selected from a range between 0.001 nM-100  $\mu\text{M}$ . Compounds were assayed in Millipore MHVB multiscreen plates (Cat# MHVBN4550) by making the following sequential additions to duplicate wells: (i) 100  $\mu\text{l}$ /well of binding buffer containing 1.5 mM  $\text{MnCl}_2$ ; (ii) 10  $\mu\text{l}$ /well  $^{125}\text{I}$ -VCAM-Ig in binding buffer (final assay concentration < 500 pM); (iii) 1.5  $\mu\text{l}$ /well test compound or DMSO alone; (iv) 38  $\mu\text{l}$ /well RPMI-8866 cell

30


suspension ( $1.25 \times 10^6$  cells/well). The plates were incubated at room temperature for 45 minutes on a plate shaker at 200 rpm, filtered on a vacuum box, and washed on the same apparatus by the addition of 100  $\mu\text{L}$  of binding buffer containing 1 mM  $\text{MnCl}_2$ . After insertion of the

5 multiscreen plates into adapter plates (Packard, Meriden, CT, cat# 6005178), 100  $\mu\text{L}$  of Microscint-20 (Packard cat# 6013621) was added to each well. The plates were then sealed, placed on a shaker for 30 seconds, and counted on a Topcount microplate scintillation counter (Packard). Control wells containing DMSO alone were used to

10 determine the level of VCAM-Ig binding corresponding to 0% inhibition. Wells in which cells were omitted were used to determine the level of binding corresponding to 100% inhibition. Percent inhibition was then calculated for each test well and the  $\text{IC}_{50}$  was determined from a ten point titration using a validated four parameter fit algorithm.

## WHAT IS CLAIMED IS:

1. A method for the treatment of diseases, disorders, conditions or symptoms mediated by cell adhesion in a mammal which comprises administering to said mammal an effective amount of a compound of Formula I:



- 10 or a pharmaceutically acceptable salt thereof wherein:

- R<sup>1</sup> is
- 1) C<sub>1</sub>-10alkyl,
  - 2) C<sub>2</sub>-10alkenyl,
  - 3) C<sub>2</sub>-10alkynyl,
  - 4) Cy,
  - 5) Cy-C<sub>1</sub>-10alkyl,
  - 6) Cy-C<sub>2</sub>-10alkenyl,
  - 7) Cy-C<sub>2</sub>-10alkynyl,

- wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from R<sup>a</sup>; and Cy is optionally substituted with one to four substituents independently selected from R<sup>b</sup>;

R<sup>2</sup> and R<sup>3</sup> are independently selected from

- 1) hydrogen, or
- 2) a group selected from R<sup>1</sup>;

25

- R<sup>4</sup> is
- 1) hydrogen,
  - 2) C<sub>1</sub>-10alkyl,
  - 3) C<sub>2</sub>-10alkenyl,
  - 4) C<sub>2</sub>-10alkynyl,

30

- 5) aryl,
- 6) aryl-C<sub>1</sub>-10alkyl,

- 7) heteroaryl,
- 8) heteroaryl-C<sub>1-10</sub>alkyl,

wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to four substituents independently selected from R<sup>a</sup>; and aryl and

- 5 heteroaryl optionally substituted with one to four substituents independently selected from R<sup>b</sup>; or

10 R<sup>3</sup>, R<sup>4</sup> and the carbon to which they are attached form a 3-7 membered mono- or bicyclic ring containing 0-2 heteroatoms selected from N, O and S;

- R<sup>5</sup> is
- 1) hydrogen,
  - 2) C<sub>1-10</sub> alkyl,
  - 3) Cy, or
  - 15 4) Cy-C<sub>1-10</sub> alkyl,

wherein alkyl is optionally substituted with one to four substituents independently selected from R<sup>a</sup>; and Cy is optionally substituted with one to four substituents independently selected from R<sup>b</sup>;

- 20 R<sup>6</sup> is
- 1) hydrogen,
  - 2) C<sub>1-10</sub>alkyl,
  - 3) C<sub>2-10</sub>alkenyl,
  - 4) C<sub>2-10</sub>alkynyl,
  - 5) Cy,
  - 25 6) Cy-C<sub>1-10</sub>alkyl,
  - 7) Cy-C<sub>2-10</sub>alkenyl,
  - 8) Cy-C<sub>2-10</sub>alkynyl,

wherein alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents selected from R<sup>x</sup>, and Cy is optionally substituted with

30 one to four substituents independently selected from R<sup>y</sup>;

- R<sup>7</sup> is
- 1) hydrogen,
  - 2) C<sub>1-10</sub>alkyl,

- 3) C<sub>2-10</sub>alkenyl,
- 4) C<sub>2-10</sub>alkynyl,
- 5) aryl,
- 6) aryl-C<sub>1-10</sub>alkyl,
- 5 7) heteroaryl,
- 8) heteroaryl-C<sub>1-10</sub>alkyl,

wherein alkyl, alkenyl and alkynyl are optionally substituted with one to four substituents selected from R<sub>x</sub>, and aryl and heteroaryl are optionally substituted with one to four substituents independently selected from R<sub>y</sub>; or

R<sub>6</sub>, R<sub>7</sub> together with the atoms to which they are attached form a 3-7 membered ring containing 0-2 heteroatoms selected from N, O and S;

- 15 R<sup>a</sup> is
- 1) Cy, or
  - 2) a group selected from R<sub>x</sub>;
- wherein Cy is optionally substituted with one to four substituents independently selected from R<sup>c</sup>;

- 20 R<sup>b</sup> is
- 1) a group selected from R<sup>a</sup>,
  - 2) C<sub>1-10</sub> alkyl,
  - 3) C<sub>2-10</sub> alkenyl,
  - 4) C<sub>2-10</sub> alkynyl,
  - 5) aryl C<sub>1-10</sub>alkyl,
  - 25 6) heteroaryl C<sub>1-10</sub> alkyl,

wherein alkyl, alkenyl, alkynyl, aryl, heteroaryl are optionally substituted with a group independently selected from R<sup>c</sup>;

- 30 R<sup>c</sup> is
- 1) halogen,
  - 2) amino,
  - 3) carboxy,
  - 4) C<sub>1-4</sub>alkyl,
  - 5) C<sub>1-4</sub>alkoxy,

- 6) aryl,
- 7) aryl C<sub>1-4</sub>alkyl, or
- 8) aryloxy;

- 5 R<sup>d</sup> and R<sup>e</sup> are independently selected from hydrogen, C<sub>1-10</sub>alkyl, C<sub>2-10</sub>alkenyl, C<sub>2-10</sub>alkynyl, Cy and Cy C<sub>1-10</sub>alkyl, wherein alkyl, alkenyl, alkynyl and Cy is optionally substituted with one to four substituents independently selected from R<sup>c</sup>; or
- 10 R<sup>d</sup> and R<sup>e</sup> together with the atoms to which they are attached form a heterocyclic ring of 5 to 7 members containing 0-2 additional heteroatoms independently selected from oxygen, sulfur and nitrogen;

R<sup>f</sup> and R<sup>g</sup> are independently selected from hydrogen, C<sub>1-10</sub>alkyl, Cy and Cy C<sub>1-10</sub>alkyl; or

- 15 R<sup>f</sup> and R<sup>g</sup> together with the carbon to which they are attached form a ring of 5 to 7 members containing 0-2 heteroatoms independently selected from oxygen, sulfur and nitrogen;

- R<sup>h</sup> is
- 1) hydrogen,
  - 20 2) C<sub>1-10</sub>alkyl,
  - 3) C<sub>2-10</sub>alkenyl,
  - 4) C<sub>2-10</sub>alkynyl,
  - 5) cyano,
  - 6) aryl,
  - 25 7) aryl C<sub>1-10</sub>alkyl,
  - 8) heteroaryl,
  - 9) heteroaryl C<sub>1-10</sub>alkyl, or
  - 10) -SO<sub>2</sub>R<sup>i</sup>;

- wherein alkyl, alkenyl, and alkynyl are optionally substituted with one to
- 30 four substituents independently selected from R<sup>a</sup>; and aryl and heteroaryl are each optionally substituted with one to four substituents independently selected from R<sup>b</sup>;

- $R^i$  is
- 1) C<sub>1-10</sub>alkyl,
  - 2) C<sub>2-10</sub>alkenyl,
  - 3) C<sub>2-10</sub>alkynyl, or
  - 4) aryl;
- 5 wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from  $R^c$ ;
- $R^x$  is
- 1)  $-OR^d$ ,
  - 2)  $-NO_2$ ,
  - 10 3) halogen
  - 4)  $-S(O)_mR^d$ ,
  - 5)  $-SR^d$ ,
  - 6)  $-S(O)_2OR^d$ ,
  - 7)  $-S(O)_mNR^dR^e$ ,
  - 15 8)  $-NR^dR^e$ ,
  - 9)  $-O(CR^fR^g)_nNR^dR^e$ ,
  - 10)  $-C(O)R^d$ ,
  - 11)  $-CO_2R^d$ ,
  - 12)  $-CO_2(CR^fR^g)_nCONR^dR^e$ ,
  - 20 13)  $-OC(O)R^d$ ,
  - 14)  $-CN$ ,
  - 15)  $-C(O)NR^dR^e$ ,
  - 16)  $-NR^dC(O)R^e$ ,
  - 17)  $-OC(O)NR^dR^e$ ,
  - 25 18)  $-NR^dC(O)OR^e$ ,
  - 19)  $-NR^dC(O)NR^dR^e$ ,
  - 20)  $-CR^d(N-OR^e)$ , or
  - 21)  $-CF_3$ ;
- 30  $R^y$  is
- 1) a group selected from  $R^x$ ,
  - 2) C<sub>1-10</sub> alkyl,
  - 3) C<sub>2-10</sub> alkenyl,
  - 4) C<sub>2-10</sub> alkynyl,

- 5) aryl C<sub>1-10</sub>alkyl,
- 6) heteroaryl C<sub>1-10</sub> alkyl,
- 7) cycloalkyl,
- 8) heterocyclyl;

5 wherein alkyl, alkenyl, alkynyl and aryl are each optionally substituted with one to four substituents independently selected from R<sup>x</sup>;  
Cy is cycloalkyl, heterocyclyl, aryl, or heteroaryl;

m is an integer from 1 to 2;

10

n is an integer from 1 to 10;

- X is
- 1) -C(O)OR<sup>d</sup>,
  - 2) -P(O)<sub>2</sub>OR<sup>d</sup>,
  - 15 3) -S(O)<sub>m</sub>OR<sup>d</sup>,
  - 4) -C(O)NR<sup>d</sup>R<sup>h</sup>, or
  - 5) -5-tetrazolyl.

2. A method of Claim 1 wherein said disease is selected  
20 from asthma, allergic rhinitis, multiple sclerosis, inflammation, inflammatory bowel disease and atherosclerosis.

3. A method of Claim 1 wherein said cell adhesion is mediated by VLA-4.  
25

4. A method of Claim 1 wherein in compounds of

Formula I

- R<sup>1</sup> is aryl optionally substituted with one to four substituents selected from R<sup>b</sup>;
- 30 R<sup>2</sup> is H or C<sub>1-3</sub>alkyl;
- one of R<sup>3</sup> and R<sup>4</sup> is hydrogen and the other is Cy, Cy-C<sub>1-10</sub>alkyl, or C<sub>1-10</sub>alkyl, each of which is optionally substituted as provided in Claim 1;




one of R<sup>6</sup> and R<sup>7</sup> is hydrogen and the other is aryl, aryl-C<sub>1-10</sub>alkyl, or C<sub>1-10</sub>alkyl, wherein aryl is optionally substituted with one to four substituents independently selected from R<sub>Y</sub>, and alkyl is optionally substituted with one to four substituents selected from R<sub>X</sub>

X is -C(O)OR<sup>d</sup>;

R<sup>b</sup>, R<sup>d</sup>, R<sub>X</sub> and R<sub>Y</sub> are as defined in Claim 1.

5. A compound having the formula Ia:



Ia

or a pharmaceutically acceptable salt thereof,

wherein

R<sup>2</sup> is 1) hydrogen, or  
2) C<sub>1-6</sub>alkyl;

one of R<sup>3</sup> and R<sup>4</sup> is hydrogen and the other is

1) aryl,  
2) heteroaryl,  
3) aryl-C<sub>1-10</sub>alkyl,  
4) heteroaryl-C<sub>1-10</sub>alkyl, or  
5) C<sub>1-10</sub>alkyl,

wherein aryl and heteroaryl are each optionally substituted with one to four substituents selected from R<sup>b</sup>; and alkyl is optionally substituted with one to four substituents independently selected from R<sup>a</sup>; or R<sup>3</sup>, R<sup>4</sup> and the carbon to which they are attached form a 3-7 membered mono- or bicyclic ring containing 0-2 heteroatoms selected from N, O and S;

one of R<sup>6</sup> and R<sup>7</sup> is hydrogen and the other is

- 1) aryl,
- 2) aryl-C<sub>1-10</sub>alkyl, or
- 3) C<sub>1-10</sub>alkyl,

- 5 wherein aryl is optionally substituted with one to four substituents independently selected from R<sup>Y</sup>, and alkyl is optionally substituted with one to four substituents selected from R<sup>X</sup>; and R<sup>a</sup>, R<sup>b</sup>, R<sup>X</sup> and R<sup>Y</sup> are as defined in Claim 1.

- 10 6. A compound of Claim 5 wherein R<sup>b</sup> is selected from C<sub>1-10</sub>alkoxy, halogen, cyano, and trifluoromethyl; R<sup>2</sup> is H or methyl; one of R<sup>3</sup> and R<sup>4</sup> is hydrogen and the other is phenyl, Cy-C<sub>1-3</sub>alkyl (wherein Cy is thienyl, pyridyl or phenyl optionally substituted with one or two groups selected from phenyl, hydroxy, halogen, cyano, and nitro);  
15 and C<sub>1-6</sub>alkyl optionally substituted with CO<sub>2</sub>H or CONH<sub>2</sub>; one of R<sup>6</sup> and R<sup>7</sup> is hydrogen and the other is aryl-C<sub>1-2</sub>alkyl, wherein aryl is optionally substituted with one to two substituents independently selected from halogen; and  
20 R<sup>Y</sup> is as defined in Claim 1.

7. A compound selected from the group consisting of:  
N-(3,4-dimethoxybenzenesulfonyl)-(L)-3-(2-thienyl)alanyl-norleucine;  
N-(3,4-dimethoxybenzenesulfonyl)-(L)-3,3-diphenylalanyl-(L)-norleucine;  
25 N-(3,4-dimethoxybenzenesulfonyl)-(L)-aspartyl-(L)-norleucine;  
N-(3,4-dimethoxybenzenesulfonyl)-(L)-3-(3-pyridyl)alanyl-(L)-norleucine;  
N-(3,4-dimethoxybenzenesulfonyl)-(L)-alanyl-(L)-norleucine;  
N-(3,4-dimethoxybenzenesulfonyl)-(L)-homophenylalanyl-(L)-norleucine;  
N-(3,4-dimethoxybenzenesulfonyl)-(L)-valyl-(L)-3-(2-naphthyl)alanine;  
30 N-(3,4-dimethoxybenzenesulfonyl)-(L)-asparagyl-(L)-norleucine;  
N-(3,4-dimethoxybenzenesulfonyl)-(L)-isoleucyl-(L)-3-(2-naphthyl)alanine;

- N-(3,4-dimethoxybenzenesulfonyl)-(L)-3-(4-biphenyl)alanyl-(L)-norleucine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-N-methylvalyl-(L)-3-(2-naphthyl)alanine;  
5 N-(3,5-di(trifluoromethyl)-benzenesulfonyl)-(L)-N-methylvalyl-(L)-3-(2-naphthyl)alanine;  
N-(3,5-dichlorobenzenesulfonyl)-2(S)-aminobutyryl-(L)-3-(2-naphthyl)alanine;  
N-(3,4-dimethoxybenzenesulfonyl)-(L)-N-methylisoleucyl-(L)-3-(2-naphthyl)alanine;  
10 N-(3,4-dimethoxybenzenesulfonyl)-(L)-N-methylvalyl-(L)-3-(2-naphthyl)alanine;  
N-(3,4-dimethoxybenzenesulfonyl)-(L)-sarcosinyl-(L)-3-(2-naphthyl)alanine;  
15 N-(3,4-dimethoxybenzenesulfonyl)-(L)-phenylalanyl-(L)-norleucine;  
N-(3,4-dimethoxybenzenesulfonyl)-(L)-N-methylphenylalanyl-(L)-norleucine;  
N-(3,4-dimethoxybenzenesulfonyl)-(L)- $\alpha$ -phenylglycyl-(L)-norleucine;  
N-(3,4-dimethoxybenzenesulfonyl)-(L)-N-methylalanyl-(L)-norleucine;  
20 N-(3,5-dichlorobenzenesulfonyl)-(L)-tyrosyl-(L)-4-fluorophenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-3-(4-pyridyl)alanine-(L)-4-fluorophenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-phenylalanyl-(L)-4-fluorophenylalanine;  
25 N-(3,5-dichlorobenzenesulfonyl)-(L)-4-fluorophenylalanyl-(L)-4-fluorophenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-3,5-difluorophenylalanyl-(L)-4-fluorophenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-N-methylaspartyl-(L)-2-naphthylalanine;  
30 N-(3,5-dichlorobenzenesulfonyl)-(L)-3-(2-pyridyl)alanyl-(L)-4-fluorophenylalanine;

- N-(3,5-dichlorobenzenesulfonyl)-(L)-N-methylvalyl-(L)-3-(2-naphthyl)alanine;  
N-(3-fluorobenzenesulfonyl)-(L)-N-methylvalyl-(L)-3-(2-naphthyl)alanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-4-chlorophenylalanyl-(L)-4-  
5 fluorophenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-4-cyanophenylalanyl-(L)-4-fluorophenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-4-nitrophenylalanyl-(L)-4-fluorophenylalanine;  
10 N-(3,5-dichlorobenzenesulfonyl)-(L)-aspartyl-(L)-4-fluorophenylalanine;  
N-(3,5-dichlorobenzenesulfonyl)-(L)-3-(3-pyridyl)alanyl-(L)-4-fluorophenylalanine; and  
N-(3,5-dichlorobenzenesulfonyl)-(D)-N-methylvalyl-(L)-4-fluorophenylalanine;  
15 N-(3,5-dichlorobenzenesulfonyl)-(L)-N-methylvalyl-(L)-3-(2-naphthyl)alanine;  
N-(3-chlorobenzenesulfonyl)-2-amino-indan-2-carbonyl-(L)-4-fluorophenylalanine;  
N-(3-fluorobenzenesulfonyl)-(L)-2-phenylglycyl-(L)-4-  
20 fluorophenylalanine; and  
N-(3,5-dichlorobenzenesulfonyl)-(L)-leucyl-(L)-aspartic acid.

8. A method for the treatment of diseases, disorders, conditions or symptoms mediated by cell adhesion in a mammal which  
25 comprises administering to said mammal an effective amount of a compound of Claim 5.

9. A method for the treatment of asthma, allergic rhinitis, multiple sclerosis, atherosclerosis inflammatory bowel  
30 disorder, or inflammation in a mammal which comprises administering to said mammal an effective amount of a compound of Claim 5.

10. A pharmaceutical composition which comprises a compound of Claim 5 and a pharmaceutically acceptable carrier thereof.

## INTERNATIONAL SEARCH REPORT

 International application No.  
 PCT/US98/10952

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(6) : Please See Extra Sheet. US CL : Please See Extra Sheet. According to International Patent Classification (IPC) or to both national classification and IPC				
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAS ONLINE				
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>				
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
A, P	US 5,663,297 A (ALIG et al.) 02 September 1997, Col. 8-26.	1-10		
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.				
<table border="0"> <tr> <td>           * Special categories of cited documents:            "A" document defining the general state of the art which is not considered to be of particular relevance            "B" earlier document published on or after the international filing date            "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)            "O" document referring to an oral disclosure, use, exhibition or other means            "P" document published prior to the international filing date but later than the priority date claimed         </td> <td>           "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention            "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone            "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art            "Z" document member of the same patent family         </td> </tr> </table>			* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "B" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "Z" document member of the same patent family
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "B" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "Z" document member of the same patent family			
Date of the actual completion of the international search 17 JULY 1998		Date of mailing of the international search report 08 SEP 1998		
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer TADEO A. SOLOLA Telephone No. (703) 308-1235		

INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/10952

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

A61K 31/42, 31/415, 31/425, 31/505; C07C 255/03; C07D 207/08, 217/18, 231/04, 263/04, 277/04, 285/12, 295/023, 413/10, 417/10


A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

514/187, 208, 230.5, 255, 256, 269, 363, 373, 382, 396, 402; 546/139; 548/136, 146, 215, 241, 252, 283, 356.1, 427, 579; 558/411

- 6 144 mo f.f.

update




- 9 transfer 211 Ques.

10. Traveller's expenses  
+ Dancer  
→ ~~do not~~ ~~cover~~ expenses until  
approved by David Reiter

2) pre-synthesized

2) partially equilibrated

3) Bumpers



175 - list of parts  
instrument designed

17C - tomorrow

~~142~~ - non-specific binding columns (new case).